

## Invisalign® informed consent and agreement

### Patient's informed consent and agreement regarding Invisalign® orthodontic treatment

Your doctor has recommended the Invisalign system for your orthodontic treatment. Although orthodontic treatment can lead to a healthier and more attractive smile, you should also be aware that any orthodontic treatment (including orthodontic treatment with Invisalign aligners) has limitations and potential risks that you should consider before undergoing treatment.

### Device Description.

Invisalign aligners, developed by Align Technology, Inc. ("Align") consist of a series of clear plastic, removable appliances that move your teeth in small increments. Invisalign's product combines your doctor's diagnosis and prescription with sophisticated computer graphics technology to develop a treatment plan which specifies the desired movements of your teeth during the course of your treatment. Upon approval of a treatment plan developed by your doctor, a series of customized Invisalign aligners is produced specifically for your treatment.

### Procedure

You may undergo a routine orthodontic pre-treatment examination including radiographs (x-rays) and photographs. Your doctor will take impressions or intra-oral scans of your teeth and send them along with a prescription to the Align laboratory. Align technicians will follow your doctor's prescription to create a ClinCheck® software model of your prescribed treatment. Upon approval of the ClinCheck treatment plan by your doctor, Align will produce and ship a series of customized aligners to your doctor. The total number of aligners will vary depending on the complexity of your malocclusion and the doctor's treatment plan. The aligners will be individually numbered and will be dispensed to you by your doctor with specific instructions for use. Unless otherwise instructed by your doctor, you should wear your aligners for approximately 20 to 22 hours per day, removing them only to eat, brush and floss. As directed by your doctor, you will switch to the next aligner in the series every two weeks or as directed by your doctor. Treatment duration varies depending on the complexity of your doctor's prescription. Unless instructed otherwise, you should follow up with your doctor at a minimum of every 6 to 8 weeks.

Some patients may require bonded aesthetic attachments and/or the use of elastics during treatment to facilitate specific orthodontic movements. Patients may require additional impressions or intra-oral scans and/or refinement aligners after the initial series of aligners.

### Benefits

- Invisalign aligners offer an aesthetic alternative to conventional braces.
- Aligners are nearly invisible so many people won't realize you are in treatment.
- Treatment plans can be visualized through the ClinCheck software.
- Aligners allow for normal brushing and flossing tasks that are generally impaired by conventional braces.
- Aligners do not have the metal wires or brackets associated with conventional braces.
- The wearing of aligners may improve oral hygiene habits during treatment.
- Invisalign patients may notice improved periodontal (gum) health during treatment.

### Risks and Inconveniences

Like other orthodontic treatments, the use of Invisalign product(s) may involve some of the risks outlined below:

- (i) Failure to wear the appliances for the required number of hours per day, not using the product as directed by your doctor, missing appointments, and erupting or atypically shaped teeth can lengthen the treatment time and affect the ability to achieve the desired results;
- (ii) Dental tenderness may be experienced after switching to the next aligner in the series;
- (iii) Gums, cheeks and lips may be scratched or irritated;
- (iv) Teeth may shift position after treatment. Consistent wearing of retainers at the end of treatment should reduce this tendency;
- (v) Tooth decay, periodontal disease, inflammation of the gums or permanent markings (e.g. decalcification) may occur if patients consume foods or beverages containing sugar, do not brush and floss their teeth properly before wearing the Invisalign products, or do

## Invisalign® informed consent and agreement

not use proper oral hygiene and preventative maintenance;

- (vi) The aligners may temporarily affect speech and may result in a lisp, although any speech impediment caused by the Invisalign® products should disappear within one or two weeks;
- (vii) Aligners may cause a temporary increase in salivation or mouth dryness and certain medications can heighten this effect;
- (viii) Attachments are tooth-colored “buttons” that may be bonded to one or more teeth during the course of treatment to facilitate tooth movement and/or appliance retention. Use of attachments may make it more noticeable that you are in treatment. These will be removed after treatment is completed;
- (ix) Attachments may fall off and require replacement.
- (x) Teeth may require interproximal recontouring or slenderizing in order to create space needed for dental alignment to occur;
- (xi) The bite may change throughout the course of treatment and may result in temporary patient discomfort.
- (xii) In rare instances, slight superficial wear of the aligner may occur where patients may be grinding their teeth or where the teeth may be rubbing and is generally not a problem as overall aligner integrity and strength remain intact.
- (xiii) At the end of orthodontic treatment, the bite may require adjustment (“occlusal adjustment”).
- (xiv) Atypically-shaped, erupting, and/or missing teeth may affect aligner adaptation and may affect the ability to achieve the desired results.
- (xv) Treatment of severe open bite, severe overjet, mixed dentition, and/or skeletally narrow jaw may require supplemental treatment in addition to aligner treatment.
- (xvi) Supplemental orthodontic treatment, including the use of bonded buttons, orthodontic elastics, auxiliary appliances/ dental devices (e.g. temporary anchorage devices, sectional fixed appliances), and/or restorative dental procedures may be needed for more complicated treatment plans where aligners alone may not be adequate to achieve the desired outcome.
- (xvii) Teeth which have been overlapped for long periods of time may be missing the gingival tissue below the interproximal contact once the teeth are aligned, leading to the appearance of a “black triangle” space.
- (xviii) Aligners are not effective in the movement of dental implants.
- (xix) General medical conditions and use of medications can affect orthodontic treatment;
- (xx) Health of the bone and gums which support the teeth may be impaired or aggravated;
- (xxi) Oral surgery may be necessary to correct crowding or severe jaw imbalances that are present prior to wearing the Invisalign product. If oral surgery is required, risks associated with anesthesia and proper healing must be taken into account prior to treatment;
- (xxii) A tooth that has been previously traumatized, or significantly restored may be aggravated. In rare instances the useful life of the tooth may be reduced, the tooth may require additional dental treatment such as endodontic and/or additional restorative work and the tooth may be lost;
- (xxiii) Existing dental restorations (e.g. crowns) may become dislodged and require re-cementation or in some instances, replacement;
- (xxiv) Short clinical crowns can pose appliance retention issues and inhibit tooth movement;
- (xxv) The length of the roots of the teeth may be shortened during orthodontic treatment and may become a threat to the useful life of teeth;
- (xxvi) Product breakage is more likely in patients with severe crowding and/or multiple missing teeth;
- (xxvii) Orthodontic appliances or parts thereof may be accidentally swallowed or aspirated;
- (xxviii) In rare instances, problems may also occur in the jaw joint, causing joint pain, headaches or ear problems;
- (xxix) Allergic reactions may occur;
- (xxx) Teeth that are not at least partially covered by the aligner may undergo supraeruption;
- (xxxi) In rare instances patients with hereditary angioedema (HAE), a genetic disorder, may experience rapid local swelling of subcutaneous tissues including the larynx. HAE may be triggered by mild stimuli including dental procedures.

## Invisalign® informed consent and agreement

### Privacy Notice

The use of the Invisalign system requires that your doctor pass your personal information, including personal health information, to Align Technology, Inc. headquartered in San Jose, California (“Align”) for the purposes of processing your personal data to provide services to your doctor at his or her request.

Your doctor is the controller of any personal information that is provided to Align. Align may use and store your personal information outside your country of residence, and subject to the laws of that other jurisdiction.

Your doctor and Align will only share your personal information with other third parties if permitted by the applicable law, such as if necessary to protect your vital interests or those of another person or if necessary for the establishment, exercise or defence of legal proceedings.

Your doctor and Align will each retain your personal information for only as long as it has an ongoing legitimate need to do so, for example, to provide you with treatment and to comply with regulatory obligations. Once no longer needed, it will then be either deleted or anonymised or, if this is not possible (for example, because your personal information has been stored in backup archives), then it will be securely stored until deletion is possible.

Your doctor or Align may de-identify your personal information to use for research and educational purposes.

### Informed Consent

I have been informed by my doctor of the nature of the procedure and I understand the benefits, risks, and alternatives for an iTero intraoral scan. I have been sufficiently informed and have had the opportunity to ask questions and discuss concerns about my dental and/or orthodontic treatment with my doctor from whom I intend to receive treatment. I hereby consent to my doctor’s use of the iTero intraoral scanner as part of my dental and/or orthodontic treatment.

Due to the fact that orthodontics is not an exact science, I acknowledge that my doctor and Align have not and cannot make any guarantees or assurances concerning the outcome of my treatment. I understand that Align is not a provider of medical, dental or health care services and does not and cannot practice medicine, dentistry or give medical advice. No assurances or guarantees of any kind have been made to me by my doctor or Align, or their representatives concerning any specific outcome of my treatment.

I understand and consent to my doctor collecting and using my personal information, including, but not limited to, radiographs (x-rays), reports, charts, medical history, photographs, findings, plaster models, impressions of teeth, or intra-oral scans, prescriptions, diagnosis, medical testing, test results, billing, and other treatment records to the extent necessary for treatment, customer service, and billing – including providing it to Align as set out in the above “Privacy Notice”.

A copy of this Consent shall be considered as effective and valid as an original. I have read, understand and agree to the terms set forth in this Consent as indicated by my signature below.

Signature \_\_\_\_\_

Print Name \_\_\_\_\_

Address \_\_\_\_\_

City \_\_\_\_\_

Date \_\_\_\_\_

Witness \_\_\_\_\_

Print Name \_\_\_\_\_

Signature of Parent/Guardian \_\_\_\_\_

If the Patient lacks the legal capacity to consent, the parent or guardian must also sign this Agreement.

Align Technology, Inc.  
Align Tech Systems Canada ULC  
240 Richmond Street W  
Toronto, ON  
M5V 2C5  
(888) 822-5446  
[privacy@aligntech.com](mailto:privacy@aligntech.com)  
[www.invisalign.ca](http://www.invisalign.ca)

© 2019 Align Technology, Inc. All Rights Reserved.  
iTero®, among others, are trademarks and/or service marks of Align Technology, Inc. or one of its subsidiaries or affiliated companies and may be registered in the U.S. and/or other countries.